

Citation for published version:

Williamson, B, Blondel, P, Waggitt, J, Fraser, S, Hall, C, Armstrong, E, Bell, P & Scott, B 2015, 'Using the FLOWBEC seabed frame to understand underwater interactions between diving seabirds, prey, hydrodynamics and MREDs', Marine Renewable Energy Knowledge Exchange Programme, UK United Kingdom, 1/03/15.

Publication date:
2015

Document Version
Early version, also known as pre-print

[Link to publication](#)

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Using the FLOWBEC seabed frame to understand underwater interactions between diving seabirds, prey, hydrodynamics and MREDs

Benjamin Williamson, Philippe Blondel,
James Waggitt, Shaun Fraser,
Chris Hall, Eric Armstrong,
Paul Bell, Beth Scott

NE/J004308/1, NE/J004200/1, NE/J004332/1


NERC MREKE Internship

Innovate UK KTP / MeyGen Ltd.

b.williamson@abdn.ac.uk

Key:

- Offshore Wind - Scottish site
- Offshore Wind - Round 3 Zones
- Robin Rigg Windfarm
- Saltire Prize Regional Locational Guidance Areas (Tidal)
- The Crown Estate Wave Lease Areas
- Saltire Prize Regional Locational Guidance Areas (Wave)
- Leased Tidal Areas
- Wind Draft plan options
- Wave Draft Plan options
- Tidal Draft Plan options



Scotland:

100% renewable by 2020

≈67.2% as of 2014

EMEC ORKNEY[®]
The European Marine Energy Centre Ltd




MEYGEN
THE TIDE OF CHANGE IN CAITHNESS

FLOWBEC Frame

- **Entire water column** (plankton, fish, seabirds, marine mammals)
- Captures **movement, behaviour and interactions** with MREDs
- Self-contained, portable between sites
- Continuously samples spring/neap 2-week period
- Complemented by concurrent:
 - hydrodynamic model data
 - above water radar and bird observations







Marine X-band Radar (Bell and McCann)

- Sea-surface currents and roughness
- Target tracking (birds, mammals)
- Live stream noc.ac.uk/project/flowbec

Track speeds (m/s), 14/06/2012 11:00am




Simrad EK60 echosounder (38, 120, 200 kHz)

- bird and fish abundance, school behaviour
- multi-frequency target identification
- morphology of turbulence, plankton

Imagenex multibeam sonar (260 kHz)

- interactions of fish, diving seabirds, marine mammals with MREDs
- target tracking, evasion behaviour




ADV


- flow, turbulence, temperature, water height

Fluorometer


- chlorophyll (phytoplankton)
- turbidity



Multibeam for target tracking




EK60 for multifrequency ID








Acoustic classification ground truthed by shore observations

Multibeam sonar tracking of diving guillemots/razorbills feeding beneath a fish shoal at a wave energy site




Green = Turbine structure, **Dashed** = Expected blade radius








Multibeam sonar 3D seal and prey tracking at tidal passes



Co-registered video and sonar tracking (above)
with concurrent prey, turbulence and photo ID (right)






Concurrent video/sonar of
predator-prey interactions
and ecologically-relevant
turbulent features

Turbulence mask and parameterisation:



- excluded from biological data, useful as a covariate

School detection:


- quantitative parameterisation of biological targets




Green = Turbine structure, **Dashed** = Expected blade radius




- **Target detection** using the multibeam and EK60
- **Target tracking** using the multibeam
- **Multifrequency analysis** using the EK60



All tracked targets (mammals, birds, fish schools, individual fish) next to Atlantis turbine structure = 3909 tracks over 2 week period




Vertical overlap with turbine height = 227 tracks over 2 week period




Co-registered MBES (behaviour and turbine interaction) and EK60 (multi-frequency and turbulence metrics)


Turbine Structure



Control Site (no turbine)




Turbine Structure



Period in turbine
structure wake

Control Site (no turbine)



- Comparable hydrodynamic conditions
- Velocity deficit when in structure wake (ADV)
- But TKE (typically correlated with velocity) is higher in wake
- Suggests different mechanism (smaller-scale intense turbulence rather than natural larger-scale turbulence)

FLOWBEC frame (single point,
temporal persistence)


complemented by

RESPONSE boat surveys
(entire site, temporal
snapshot)

Waggitt *et al.* (2014) EIMR

**Suitable Physical
Conditions**

**Increased Prey
Availability**



**Presence of
Foraging Seabirds**



Investigating the ecological effects of installing and operating MREDs


- Investigate collision risk probabilities
- Define vertical habitat use and any changes in habitat use pre & post installation for a range of species
- Increase overall environmental understanding of mobile animal use of high energy sites
- Inform marine spatial planning, device design, licensing and operation
- Guide scaling-up to arrays and new site selection
- Increase predictive power to eventually reduce monitoring



Environmental Monitoring at MeyGen (UK)

Staged consent:


- 4 turbines, construction started
- then 61 turbines (86 MW)
- eventually 398 MW



Environmental Monitoring at MeyGen (UK)

Intelligent triggering of instruments across **multiple scales**

- Combines **large-scale** with **fine-detail**
- Reduces data processing / archival
- Cycle passive / active acoustics
- Trigger camera for ID / detail



Williamson, B.J. *et al.* 2015. A Self-Contained Subsea Platform for Acoustic Monitoring of the Environment Around Marine Renewable Energy Devices – Field Deployments at Wave and Tidal Energy Sites in Orkney, Scotland. *IEEE Journal of Oceanic Engineering* [10.1109/JOE.2015.2410851](https://doi.org/10.1109/JOE.2015.2410851)



NE/J004308/1, NE/J004200/1, NE/J004332/1

NERC MREKE Internship

Innovate UK KTP / MeyGen Ltd.

b.williamson@abdn.ac.uk